



Acetaminophen/Tylenol (can be given every 4-6 hours)

	6-11 lbs.	12-15 lbs.	16-21 lbs.	22-29 lbs.	30-41 lbs.	42-59 lbs.	60-71 lbs.	72-95 lbs.	Over 95 lbs.
Elixir/Children's (160mg/tsp)	--	½ tsp (2.5 ml)	¾ tsp (3.7 ml)	1 tsp (5 ml)	1 ½ tsp (1.5 ml)	2 tsp (10 ml)	2 ½ tsp (12.5 ml)	3 tsp (15 ml)	4 tsp (20 ml) (adult dose)
Suppository (80mg)	--	1 supp.	1 ½ supp.	2 supp.	3 supp.	--	--	--	--
Suppository (120mg)	--	--	1 supp.	1 ½ supp.	2 supp.	2 ½ supp.	3 supp.	--	--
Suppository (325mg)	--	--	--	--	--	--	1 supp.	1 supp.	1 ½ - 2 supp.
Tablets (325mg)	--	--	--	--	--	--	--	--	--

Ibuprofen/Motrin/Advil (can be given every 6-8 hours)

	6-11 lbs.	12-15 lbs.	16-21 lbs.	22-29 lbs.	30-41 lbs.	42-59 lbs.	60-71 lbs.	72-95 lbs.	Over 95 lbs.
Infant drops (50 mg/1.25ml)	none	> 6 months 1.25 ml	1.87 ml	2.5 ml	--	--	--	--	--
Children's Suspension (100 mg/tsp)	none	½ tsp (2.5 ml)	¾ tsp (3.7 ml)	1 tsp (5 ml)	1 ½ tsp (7.5 ml)	2 tsp (10 ml)	2 ½ tsp (12.5 ml)	3 tsp (15 ml)	4 tsp (20 ml) (adult dose)
Tablets (200 mg)	--	--	--	--	--	1 tablet	1 tablet	1 ½ tablets	2 tablets

Diphenhydramine/Benadryl for Allergic Reaction

			18-22 lbs.	23-30 lbs.	31-44 lbs.	45 lbs. or more			
Children's Benadryl 120 mg/tsp			12- 15 months ¾ tsp	16-24 months 1 tsp (5 ml)	1-6 years 1 ½ tsp (7.5 ml)	more than 6 yrs. 2 tsp (10 ml)			